

Tittle: Preparation of 4-nitrophenol

Work instructions

Task:

- 1. Make a diazotisation of 4-nitroaniline, divide the obtained solution of 4nitrobenzenediazonium sulfate into two parts (4 ml and the rest). Keep both solutions in an ice bath.
- 2. With 4 ml of the sample, carry out the demonstration of the diazonium salts by an azo copulation reaction with phenol, 2-naphthol and H-acid. Discuss the effect of structure on the colour of the compound.
- 3. The major fraction of the diazonium salt was converted by the Griess reaction to 4nitrophenol.
- 4. After crystallisation, measure the melting point and calculate the percentage yield of the reaction.

Theory

Diazotisation is the reaction of primary aromatic amines with nitrous acid to form a diazonium salt. The reaction proceeds according to the Eq:



The Griess reaction is the decomposition of a diazonium salt by hydrolysis according to Eq:



Name of the	Digitization of chemistry experiments to improve the quality and
project:	support chemistry teaching in secondary schools
Acronym:	ChemIQSoc
Project	2021-1-SK01-KA220-VET-000027995
number:	



Evidence of diazonium salt by azo copulation reactions proceeds according to Eqn:



Equipment: heating mantle, Liebig condenser, ring stand, clamps, holder, burner, tripod, wire gauze, beakers (100, 250 ml), graduated cylinder (10, 100 ml), glass rod, ice bath, scales, thermometer, pipette, pipetting balloon, test tubes, Erlenmeyer flask (250 ml), boiling chips, filter paper

Chemicals: 4-nitroaniline, sulfuric acid, sodium nitrite, phenol, 2-naphthol, H-acid, sodium hydroxide

Procedures:

1. All work is carried out with safety goggles!

Diazotisation

- 1. Pour 20 ml of water into a 200 ml beaker and carefully (SHIELD!) add 14 ml of conc. sulfuric acid, stirring constantly. To the hot solution add 7 g of finely crushed 4-nitroaniline.
- Cool the resulting solution in an ice bath to 5°C, add 25 g of ice and add a solution of 3.9 g of sodium nitrite in 18 ml of water at (0-5°C) while stirring and cooling. The diazonium salt formed makes a clear solution.
- 3. Remove 4 ml for the proof reaction, hydrolyse the rest.



Proof of diazonium salt

- 1. Divide the collected 4 ml sample into 3 tubes. Keep the temperature at 0°C.
- 2. Drop a sample of the diazonium salt successively into a solution of phenol, naphthol and H-acid (0.1 g in 3 ml of 5% NaOH). Do the same by drop reaction on filter paper.

Hydrolysis of diazonium salt

- 1. Pour 15 ml of water and 15 ml of concentrated sulfuric acid into a 250 ml flask with a reflux Liebig condenser and heat the solution to boil. Add the diazonium salt solution to the boiling solution, one at a time, using the condenser, so that the mixture continues to boil.
- 2. Rinse the residue of the diazonium salt from the beaker and the cooler with 15 ml of water. Heat the mixture to boiling for 10 minutes, then pour into a 250 ml Erlenmeyer flask, cool with water and then in an ice bath.
- 3. The eluted product is aspirated and recrystallised from hydrochloric acid diluted 1:1. Dry, calculate the yield and measure the melting point.

Chemicals	Form	H-statements	P-statements
4-Nitroaniline	Solid	H301, H301 + H311	P261, P273, P280,
		+ H331, H311,	P301 + P310, P311
		H331, H373	
H ₂ SO ₄	Liquid, 98%	H290, H314	P260, P280, P303 +
			P361 + P353, P304 +
			P340 + P310, P305 +
			P351 + P338
NaNO ₂	Solid	H272, H301, H319,	P210, P220, P221,
		H400	P301 + P330 + P331
			+ P310, P305 + P351
			+ P338, P370 + P378
Phenol	Solid, 99%	H301, H301 + H311	P260, P280, P301 +
		+ H331, H311,	P330 + P331 + P310,
		H314, H315, H319,	P303 + P361 + P353,
		H331, H341, H373,	P304 + P340 + P310,
		H411	P305 + P351 + P338
2-Naphthol	Solid, 99%	H302, H332, H400	P273
H-acid	Solid		

Management of chemical substances

Name of the	Digitization of chemistry experiments to improve the quality and
project:	support chemistry teaching in secondary schools
Acronym:	ChemIQSoc
Project	2021-1-SK01-KA220-VET-000027995
number:	



Chemicals	Form	H-statements	P-statements
NaOH	Solid	H290, H314, H315,	P260, P280, P303 +
		H319	P361 + P353, P304 +
			P340 + P310, P305 +
			P351 + P338
4-Nitrophenol	Solid	H301, H312 + H332,	P261, P301 + P310 +
(product)		H373	P330, P302 + P352 +
			P312, P304 + P340 +
			P312

Sources of risk and assessment of risk severity

There is no risk if all the principles for working with chemicals are followed and personal protective equipment (gloves, goggles, lab coat) is used.

Waste management method

These substances and their packaging must be disposed of as hazardous waste. Dispose of waste substances in a labelled container. Do not return unconsumed residues in storage bottles. Dispose of broken glass in a labelled container.

This combustible material may be incinerated in a chemical waste incinerator equipped with an afterburner and gas scrubber. Take leftover quantities and unusable solutions to a certified company for disposal. Dispose of contaminated packaging as unconsumed product.

Risk reduction measures

Wear a protective shield or goggles. Wear protective gloves. Gloves must be inspected before use. Use proper glove removal technique without touching the outer surface of the gloves to prevent skin contact with this product. Do not eat, drink or smoke while working. Wash your hands with soap and water or treat them with a protective cream when you have finished working or when you stop working. In case of an accident or if you feel unwell, inform the teacher immediately. Avoid further leakage or spillage unless there is a risk involved. Do not allow it to reach the sewer. Avoid discharging into the environment. Name of the
project:Digitization of chemistry experiments to improve the quality and
support chemistry teaching in secondary schoolsAcronym:ChemIQSocProject2021-1-SK01-KA220-VET-000027995number:ChemIQSoc



Worksheet

Calculation

- 1. Calculate the theoretical yield of 4-nitrophenol.
- 2. Calculate the practical yield of 4-nitrophenol in %.

Observation

- 1. Describe the appearance and smell of 4-nitrophenol.
- 2. Search for the basic physicochemical properties of 4-nitrophenol.

Feature	Value
Solubility in water	
Solubility in other solvents	
Melting point	

Yield and melting point

1. Record the mass of the prepared 4-nitrophenol.

 $m(C_6H_5NO_3) = \dots g$

2. Record the melting temperature of the prepared 4-nitrophenol. $t_m(C_6H_5NO_3) = \dots ^{\circ}C$

Questions

1. Characterize diazotation and describe its mechanism.



- Describe the importance of cooling the reaction mixture to 0 5 °C during diazotization. Discuss the consequences of not maintaining an accurate temperature (0 - 5 °C) during diazotation.
- 3. Compare the reactions of diazonium salt with phenol, naphthol and H-acid.
- 4. Describe why a reflux condenser is used in the hydrolysis of diazonium salt.
- 5. State what is the purpose of heating the reaction mixture.
- 6. Explain the importance of the following operations in the preparation of 4-nitrophenol.
 - a. distillation:
 - b. recrystallisation:
- 7. Consider and write what factors can affect the stability of a diazonium salt.
- 8. Describe the uses of nitrophenol.

Name of the
project:Digitization of chemistry experiments to improve the quality and
support chemistry teaching in secondary schoolsAcronym:ChemIQSocProject2021-1-SK01-KA220-VET-000027995number:ChemIQSOC



Conclusion

Briefly summarize the objective of the experiment, the main results and compare them with the expected values.

Disclaimer

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or Slovak Academic Association for International Cooperation, National Agency for the Erasmus+ Programme for Education and Training Sectors. Neither the European Union nor the granting authority can be held responsible for them.